

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 25/00	A1	(11) International Publication Number: WO 95/31100 (43) International Publication Date: 23 November 1995 (23.11.95)
<p>(21) International Application Number: PCT/US95/01283</p> <p>(22) International Filing Date: 31 January 1995 (31.01.95)</p> <p>(30) Priority Data: 08/245,743 16 May 1994 (16.05.94) US</p> <p>(71) Applicant: MCCORMAC, Dennis, J. doing business as SAFE- TEE C HEMICAL PRODUCTS COMPANY [US/US]; Suite #12-A, 4001 East Broadway, Phoenix, AZ 85040 (US).</p> <p>(72) Inventor: ABOUD, George, M.; 3939 East Glenrosa, Phoenix, AZ 85018 (US).</p> <p>(74) Agents: OGRAM, Mark, E. et al.; Ogram & Teplitz, P.C., Suite 101, 2100 North Kolb Road, Tucson, AZ 85715 (US).</p>		<p>(81) Designated States: AU, BR, CA, CN, JP, KR, NZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: ENVIRONMENTALLY SAFE INSECTICIDE</p> <p>(57) Abstract</p> <p>An environmentally-safe, broad spectrum insecticide that is toxic to invertebrate animals and non-toxic to vertebrate animals. The insecticide comprises a water solution of hydroxy acyclic acid, such as citric acid, and ionic or non-ionic surfactant thoroughly mixed together. The insecticide may be applied by aerosol, misting, spraying, or pouring on the insects or into their abodes. The composition is also an effective mosquito repellant when applied to human skin.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

ENVIRONMENTALLY SAFE INSECTICIDEBackground of the Invention:

The present invention relates generally to an environmentally safe insecticide and more particularly to novel compositions which are safe for both plant life and vertebrates in general, including human beings, but which are toxic to invertebrate pests such as insects. The composition has the further unexpected benefit of repelling mosquitos when applied to human skin.

Heretofore, insecticides used for chemically controlling insect infestation fell into five major categories namely, chlorinated hydrocarbons such as LINDANE, METHOXYCHLOR, and the like; organophosphates such as MALATHION, DIBROM, DIAZINON, PHOSPHAMIDON and the like; carbonate compounds such as BAYGON, SEVIN and the like; inorganic compounds such as, arsenic, sulphur, borax, and the like; and botanical compounds such as pyrethrin, strychnine, nicotine, and the like.

The prior chemical insecticides, excluding the botanical compounds, possess several disadvantages. Most of them are highly toxic to fish, wildlife and humans and must be used with caution. Further, the active ingredients are generally not naturally occurring biological constituents and, as such, tend to persist for long lengths of time after their initial application. Consider, for example, the chlorinated hydrocarbons (Petrochemical hydrocarbons are suspect carcinogenic agents) which because of their persistence have been shown to pass through the animal food chain and cause egg shell thinning and egg breakage in many species of birds.

Besides their initial toxic effect, these compounds have sub-acute effects on non-target fauna and flora thereby causing biochemical, behavioral and physiological changes as well as reproductive failure. Furthermore, their adverse effect on the environment in general is widely known and needs no elaboration here.

The ideal insecticide is one that adversely affects only the

target species, sublimates without residue when desired, and has no negative environmental effects. As usual, the ideal is a potentially unattainable goal which, nonetheless, is worth striving for.

5 The desiratum of the present invention was to locate and identify a substance or substances naturally occurring in some part of the environment, which has positive effects, or is at least neutral, upon all fauna and flora other than the targeted pests and is especially lethal upon those pests. Such a
10 substance must also be free of metabolites or breakdown components which themselves are toxic or cause environmental harm. The search for such a substance has been ongoing. To date, however, this search has discovered no alternatives which are truly satisfactory.

15 One prior art effort to create a non-toxic insecticide used for controlling insect infestation is based on using a non-toxic carrier such as diatomaceous earth which is thoroughly laced with a bait substance such as cane molasses, cane sugar, dextrin, and
20 the like and which after ingestion mechanically kills the insects by lacerating the body and causing the loss of vital fluids. However, such an insecticide poses a threat to non-targeted life and as such does not attain the aforestated desiratum.

25 BRIEF SUMMARY OF THE INVENTION

 The present invention relates generally to a new and novel environmentally safe insecticide which is toxic to invertebrate animals such as insects but which is completely harmless, in the
30 concentrations of usage required, to vertebrates, that is, animals with spines. Surprisingly, the essential active ingredient of the insecticide is present in every living plant and animal. Yet it was discovered that when an excess of the active ingredient is delivered to an insect, it can disrupt the
35 nervous system of the insect which, in turn, causes respiratory

failure and death.

The essential ingredients of the present invention are a hydroxy acyclic acid selected from the group consisting of citric, lactic, tartaric, malic, glycolic (gallic), tannic or ascorbic acid, and an ionic or non-ionic surfactant. The purpose of the surfactant is to reduce the surface tension of the insecticidal composition, so that when such composition is applied to the body of an insect, the penetration of the hydroxy acyclic acid into the insect's nervous system is facilitated so as to disrupt the normal respiratory function of the insect and thereby suffocate it.

The insecticide is formed by mixing a dry anhydrous acid with tap water and adding the surfactant thereto. The solution will generally contain up to about 10.5 ounces of the hydroxy acyclic acid and up to about 60 ml of the surfactant per liter of water.

The solution thus prepared and mixed can be applied using conventional delivery techniques such as with an aerosol dispenser or it can be deposited directly upon suspected insect infestation or burrowing sites as by misting, spraying or pouring.

Accordingly, a prime object of the present invention is to provide a new and useful insecticide which is environmentally safe but is selectively toxic to invertebrate animals.

Another object of the present invention is to provide a new and useful insecticide containing hydroxy acyclic acids and an ionic or non-ionic surfactant as its essential active ingredients and which is effective to destroy insects but possess no deleterious action on surrounding flora and fauna.

These and still further objects as shall hereinafter appear are readily fulfilled by the present invention in a remarkably unexpected manner as will be readily discerned from the following detailed description of an exemplary embodiment thereof.

DESCRIPTION OF PREFERRED EMBODIMENTS

The present invention relates to a new and novel environmentally safe insecticide which is toxic to invertebrate animals such as insects but which is otherwise harmless.

This invention is predicated on the use of a composition
5 containing hydroxy acid belonging to the acyclic family of compounds and an ionic or non-ionic surfactant. The specific hydroxy acid used to exemplify this invention is citric acid (2-hydroxy-1,2,3,-propanetricarboxylic acid)
H₀₀CCH₂C (OHCOOH) CH₂ COOH. H₂O, although lactic, tartaric, malic,
10 glycolic (gallic), tannic, and ascorbic acid, the so-called "food acids", can be substituted therefor with equally favorable results.

Any ionic or non-ionic surfactant may be used together with the hydroxy acyclic acid in formulating the insecticidal
15 composition of the present invention. Suitable surfactants include any of the following conventional surfactant types:

- (a) alcohols;
- (b) alkanolamides;
- (c) alkoxylated derivatives of alcohols, alkyl phenols,
20 amines, amides, fatty acids, fatty esters and oils;
- (d) amine acetates and oxides;
- (e) aryl, alkyl, aralkyl and alkaryl sulfonates and sulfates;
- (f) betaine derivatives;
- 25 (g) carboxylated alcohol ethoxylates;
- (h) esters of fatty acids, glycols, glycerols, sucrose, glucose and phosphoric acid;
- (i) fluorcarbon-based surfactants;
- (j) imidazolines, imadazolines and other heterocyclic
30 surfactants;
- (k) isethionates;
- (l) lanolin-based derivatives;
- (m) lecithin and lecithin derivatives;
- (n) lignin and lignin derivatives;
- 35 (o) polysaccharides, polyacrylates, polyacrylamides, and

- other polymeric and block polymeric surfactants;
- (p) protein-based surfactants;
 - (q) quaternary surfactants;
 - (r) sarcosine derivatives;
 - 5 (s) silicone-based surfactants;
 - (t) soaps;
 - (u) sorbitan derivatives;
 - (v) sulfates and sulfonates of amines, amides, olefins, petroleum, oils, fatty acids, fatty esters, ethoxylated
 - 10 alcohols and ethoxylated alkyl phenols;
 - (w) sulfosuccinates and sulfosuccinamates;
 - (x) taurates; and
 - (y) thio, mercapto and phosphorous derivatives.

Preferable surfactants include alkaryl polyethylene glycols, alkyl polyethylene glycols, salts of dodecylated oxydibenzene

15 disulfonate, ethoxylated nonylphenol surfactants, aliphatic polyoxyethylene ether surfactants, polyoxyethylene alkyl ether surfactants, and castor oil derivatives.

In formulating the insecticidal composition of the present invention, up to about 10.5 ounces of the hydroxy acyclic acid is

20 mixed into one (1) liter of tap water, and thereafter up to about 60 ml of the surfactant is added into the solution. The optimum amounts of these two components will vary within these broad limits, depending upon the specific hydroxy acyclic acid and surfactant employed. When using citric acid and one of the

25 surfactants listed above as preferred, typical formulations will contain from about 1.5 to about 6.5 ounces, preferably from about 1.5 to about 2.5 ounces, of the citric acid, and from about 1 to about 10 ml, preferably from about 4 to about 6 ml, of the

30 surfactant per liter of water.

The insecticide is applied by any suitable means i.e. aerosol, misting, spraying or pouring down into insect mounds or on to insect nests.

The formulation as described above is environmentally safe,

35 since all fauna and flora contain citric acid in their cells,

that is, citric acid (and other food acids) is a natural biological constituent. The insecticide function of the substance occurs in non-vertebrate animals when an excess of the acyclic hydroxy acid such as citric acid is administered thereto.

5 Respiratory failure results in the insect because the normal nervous system rhythm is disrupted by the insecticide. Of particular value is the fact that any excess insecticide (acid base) which falls on alkali soil will self-neutralize itself and be used by the surrounding plants. Likewise, excess insecticide
10 will also remain on acid soil as a residual and eventually be used by the surrounding plants.

The insecticide of the present invention totally eliminates the use of petro and chlorinated solvents for insect control and thereby precludes the contamination of ground water, streams, and
15 air with these environmentally hostile compounds.

It is further noteworthy that the hydroxy acyclic acids, such as citric acid, are hydroscopic, that is, after application, when the water evaporates therefrom, the residue of the insecticide continues to draw insects. Since citric acid is
20 derived by mold fermentation of lemons, limes, pineapple juice, molasses, and the like, it is a natural bait for insects. Upon the occurrence of high humidity or rain fall, the insecticide when disposed on a hard surface, will rehydrate itself.

A further advantage arises when citric acid is the active
25 ingredient in this insecticide, in that it is not only safe for humans but because it is consumed by humans regularly in the form of oranges, lemons, lime juice, tomatoes, and juice-flavoring extracts, confections, soft drinks, medicines, and is used as an antioxidant in many foods.

30 The insecticide of the present invention, has been extensively tested and found effective in the elimination of regular ants, fireants, cockroaches including both German and the big sewer variety, mosquitoes, locusts, flies, crickets, grasshoppers, tomato worms, beetles, spiders, earwigs
35 (dermaterpa), wasps, hornets, and yellow jackets.

This invention has applicability for residential, industrial and agricultural use.

To assist in the understanding of the present invention and not by way of limitation, the following examples are presented:

5

EXAMPLE I

To one liter of tap water, 2.1 ounces of citric acid are mixed with stirring to assure uniform dispersion of the acid throughout the water. To the diluted acid-water mixture, 5 ml of alkyl polyethylene glycol (available from Union Carbide under the brand name TERGITOL-TMN) is added with additional stirring to complete the formulation of an environmentally safe insecticide embodying the present invention.

15

EXAMPLE II

The procedure of Example I was repeated using sodium salt of dodecylated oxydibenzene disulfonate (available from Dow Chemical under the brand name BENAX 2AI) as surfactant to create an environmentally safe insecticide embodying the present invention.

20

EXAMPLE III

The procedure of Example I was repeated using an ethoxylated nonylphenol surfactant (available from Monsanto Company under the brand name STEROX-N) as the surfactant.

25

EXAMPLE IV

The preparation prepared according to Example I was placed in a spray bottle and sprayed over a nest of approximately 15 sewer cockroaches. After twenty seconds, all of the roaches were dead.

30

35

EXAMPLE V

A preparation prepared according to Example II was loaded in an atomizer and sprayed upon a nest of wasps and after ten seconds, all of the wasps were dead.

5

EXAMPLE VI

A preparation prepared according to Example III was placed in a pump bottle and sprayed at a swarm of mosquitoes. All of the mosquitoes died within five seconds.

10

* * *

A number of commercially available ionic or non-ionic surfactants can be used to prepare the present invention including alkylaryl polyethylene glycol and alkyl polyethylene glycol (available from Union Carbide under the brand name TERGITOL), the sodium salt of dodecylated oxydibenzene disulfonate (BENAX 2AI Dow); ethoxylated nonylphenol surfactants, aliphatic polyoxyethylene ether, and polyoxyethylene alkyl ether (STEROX, Monsanto); and various castor oil derivatives (SURFACTOL 13, Baker Castor Oil Co.).

Further various of the known so-called "food acids" can be used in place of the citric acid without significantly diminishing the effectiveness of the insecticide. Thus, lactic, tartaric, malic, glycolic, gallic, tannic or ascorbic acid can be substituted for the citric acid specified herein.

A further benefit of the present invention arises from its unexpected properties as an insect repellant when applied to human skin.

30

From the foregoing, it is readily apparent that a useful embodiment of the present invention has been herein described and illustrated which fulfills all of the aforestated objectives in a remarkably unexpected fashion. It is of course understood that such modifications, alterations and adaptations as may readily

35

occur to the artisan confronted with this disclosure are intended within the spirit of this disclosure which is limited only by the scope of the claims appended hereto.

What is claimed is:

1. An environmentally safe insecticide which is non-toxic to vertebrates, consisting essentially of an hydroxy acyclic acid, an ionic or non-ionic surfactant, and water, the amount of said hydroxy acyclic acid being up to about 10.5 ounces per liter of water, the amount of said surfactant being up to about 60 ml per liter of water, said hydroxy acyclic acid being selected from the group consisting of citric, lactic, tartaric, malic, glycolic, tannic and ascorbic acids, said surfactant being selected from the group consisting of:
 - (a) alcohols;
 - (b) alkanolamides;
 - (c) alkoxyated derivatives of alcohols, alkyl phenols, amines, amides, fatty acids, fatty esters and oils;
 - (d) amine acetates and oxides;
 - (e) aryl, alkyl, aralkyl and alkaryl sulfonates and sulfates;
 - (f) betaine derivatives;
 - (g) carboxylated alcohol ethoxylates;
 - (h) esters of fatty acids, glycols, glycerols, sucrose, glucose and phosphoric acid;
 - (i) fluorocarbon-based surfactants;
 - (j) imidazolines, imadazolines and other heterocyclic surfactants;
 - (k) isethionates;
 - (l) lanolin-based derivatives;
 - (m) lecithin and lecithin derivatives;
 - (n) lignin and lignin derivatives;
 - (o) polysaccharides, polyacrylates, polyacrylamides, and other polymeric and block polymeric surfactants;
 - (p) protein-based surfactants;
 - (q) quaternary surfactants;
 - (r) sarcosine derivatives;
 - (s) silicone-based surfactants;

- (t) soaps;
- (u) sorbitan derivatives;
- (v) sulfates and sulfonates of amines, amides, olefins,
petroleum, oils, fatty acids, fatty esters, ethoxylated
alcohols and ethoxylated alkyl phenols;
- (w) sulfosuccinates and sulfosuccinamates;
- (x) taurates; and
- (y) thio, mercapto and phosphorous derivatives.

2. The insecticide of claim 1, wherein said hydroxy
acyclic acid is citric acid.

3. The insecticide of claim 1, wherein said surfactant is
selected from the group consisting of alkaryl polyethylene
glycols, alkyl polyethylene glycols, salts of dodecylated
oxydibenzene disulfonate, ethoxylated nonylphenol surfactants,
aliphatic polyoxyethylene ether surfactants, polyoxyethylene
alkyl ether surfactants, and castor oil derivatives.

4. The insecticide of claim 1, wherein the amount of said
hydroxy acyclic acid is from about 1.5 to about 6.5 ounces per
liter of water.

5. The insecticide of claim 4, wherein the amount of said
hydroxy acyclic acid is from about 1.5 to about 2.5 ounces per
liter of water.

6. The insecticide of claim 1, wherein the amount of said
surfactant is from about 1 to about 10 ml per liter of water.

7. The insecticide of claim 6, wherein the amount of said
surfactant is from about 4 to about 6 ml per liter of water.

8. A method of killing insects comprising applying to said
insects or to a target site of insect infestation an

environmentally safe insecticide which is non-toxic to vertebrates and which consists essentially of an hydroxy acyclic acid, an ionic or non-ionic surfactant, and water, the amount of said hydroxy acyclic acid being up to about 10.5 ounces per liter of water, the amount of said surfactant being up to about 60 ml per liter of water, said hydroxy acyclic acid being selected from the group consisting of citric, lactic, tartaric, malic, glycolic, tannic and ascorbic acids, said surfactant being selected from the group consisting of:

- 10 (a) alcohols;
- (b) alkanolamides;
- (c) alkoxyated derivatives of alcohols, alkyl phenols, amines, amides, fatty acids, fatty esters and oils;
- (d) amine acetates and oxides;
- 15 (e) aryl, alkyl, aralkyl and alkaryl sulfonates and sulfates;
- (f) betaine derivatives;
- (g) carboxylated alcohol ethoxylates;
- (h) esters of fatty acids, glycols, glycerols, sucrose, glucose and phosphoric acid;
- 20 (i) fluorocarbon-based surfactants;
- (j) imidazolines, imadazolines and other heterocyclic surfactants;
- (k) isethionates;
- 25 (l) lanolin-based derivatives;
- (m) lecithin and lecithin derivatives;
- (n) lignin and lignin derivatives;
- (o) polysaccharides, polyacrylates, polyacrylamides, and other polymeric and block polymeric surfactants;
- 30 (p) protein-based surfactants;
- (q) quaternary surfactants;
- (r) sarcosine derivatives;
- (s) silicone-based surfactants;
- (t) soaps;
- 35 (u) sorbitan derivatives;

- (v) sulfates and sulfonates of amines, amides, olefins, petroleum, oils, fatty acids, fatty esters, ethoxylated alcohols and ethoxylated alkyl phenols;
- (w) sulfosuccinates and sulfosuccinamates;
- 5 (x) taurates; and
- (y) thio, mercapto and phosphorous derivatives.

9. The method of claim 8, wherein said hydroxy acyclic acid is citric acid.

10

10. The method of claim 8, wherein said surfactant is selected from the group consisting of alkaryl polyethylene glycols, alkyl polyethylene glycols, salts of dodecylated oxydibenzene disulfonate, ethoxylated nonylphenol surfactants, 15 aliphatic polyoxyethylene ether surfactants, polyoxyethylene alkyl ether surfactants, and castor oil derivatives.

15

11. The method of claim 8, wherein the amount of said hydroxy acyclic acid is from about 1.5 to about 6.5 ounces per 20 liter of water.

20

12. The method of claim 11, wherein the amount of said hydroxy acyclic acid is from about 1.5 to about 2.5 ounces per liter of water.

25

13. The method of claim 8, wherein the amount of said surfactant is from about 1 to about 10 ml per liter of water.

30

14. The method of claim 13, wherein the amount of said surfactant is from about 4 to about 6 ml per liter of water.

15. A method of repelling insects from the human skin comprising applying to said human skin an environmentally safe insecticide which is non-toxic to vertebrates and which consists 35 essentially of an hydroxy acyclic acid, an ionic or non-ionic

35

surfactant, and water, the amount of said hydroxy acyclic acid being up to about 10.5 ounces per liter of water, the amount of said surfactant being up to about 60 ml per liter of water, said hydroxy acyclic acid being selected from the group consisting of
5 citric, lactic, tartaric, malic, glycolic, tannic and ascorbic acids, said surfactant being selected from the group consisting of:

- (a) alcohols;
- (b) alkanolamides;
- 10 (c) alkoxyated derivatives of alcohols, alkyl phenols, amines, amides, fatty acids, fatty esters and oils;
- (d) amine acetates and oxides;
- (e) aryl, alkyl, aralkyl and alkaryl sulfonates and sulfates;
- 15 (f) betaine derivatives;
- (g) carboxylated alcohol ethoxylates;
- (h) esters of fatty acids, glycols, glycerols, sucrose, glucose and phosphoric acid;
- (i) fluorocarbon-based surfactants;
- 20 (j) imidazolines, imadazolines and other heterocyclic surfactants;
- (k) isethionates;
- (l) lanolin-based derivatives;
- (m) lecithin and lecithin derivatives;
- 25 (n) lignin and lignin derivatives;
- (o) polysaccharides, polyacrylates, polyacrylamides, and other polymeric and block polymeric surfactants;
- (p) protein-based surfactants;
- (q) quaternary surfactants;
- 30 (r) sarcosine derivatives;
- (s) silicone-based surfactants;
- (t) soaps;
- (u) sorbitan derivatives;
- (v) sulfates and sulfonates of amines, amides, olefins,
35 petroleum, oils, fatty acids, fatty esters, ethoxylated

15

- alcohols and ethoxylated alkyl phenols;
(w) sulfosuccinates and sulfosuccinamates;
(x) taurates; and
(y) thio, mercapto and phosphorous derivatives.

5

16. The method of claim 15, wherein said hydroxy acyclic acid is citric acid.

10 17. The method of claim 15, wherein said surfactant is selected from the group consisting of alkaryl polyethylene glycols, alkyl polyethylene glycols, salts of dodecylated oxydibenzene disulfonate, ethoxylated nonylphenol surfactants, aliphatic polyoxyethylene ether surfactants, polyoxyethylene alkyl ether surfactants, and castor oil derivatives.

15

18. The method of claim 15, wherein the amount of said hydroxy acyclic acid is from about 1.5 to about 6.5 ounces per liter of water.

20

19. The method of claim 18, wherein the amount of said hydroxy acyclic acid is from about 1.5 to about 2.5 ounces per liter of water.

25

20. The method of claim 15, wherein the amount of said surfactant is from about 1 to about 10 ml per liter of water.

21. The method of claim 20, wherein the amount of said surfactant is from about 4 to about 6 ml per liter of water.

30

22. An environmentally safe insecticide which is non-toxic to vertebrates, consisting essentially of an hydroxy acyclic acid, an ionic or non-ionic surfactant, and water, the amount of said hydroxy acyclic acid being up to about 10.5 ounces per liter of water, the amount of said surfactant being up to about 60 ml per liter of water, said hydroxy acyclic acid being selected from

35

the group consisting of citric, lactic, tartaric, malic, glycolic, tannic and ascorbic acids.

23. The insecticide of claim 22, wherein said hydroxy
5 acyclic acid is citric acid.

24. A method of killing insects comprising applying to said
insects or to a target site of insect infestation an
environmentally safe insecticide which is non-toxic to
10 vertebrates and which consists essentially of an hydroxy acyclic
acid, an ionic or non-ionic surfactant, and water, the amount of
said hydroxy acyclic acid being up to about 10.5 ounces per liter
of water, the amount of said surfactant being up to about 60 ml
per liter of water, said hydroxy acyclic acid being selected from
15 the group consisting of citric, lactic, tartaric, malic,
glycolic, tannic and ascorbic acids.

25. The method of claim 24, wherein said hydroxy acyclic
acid is citric acid.
20

26. A method of repelling insects from human skin
comprising applying to said human skin an environmentally safe
insecticide which is non-toxic to vertebrates and which consists
essentially of an hydroxy acyclic acid, an ionic or non-ionic
25 surfactant, and water, the amount of said hydroxy acyclic acid
being up to about 10.5 ounces per liter of water, the amount of
said surfactant being up to about 60 ml per liter of water, said
hydroxy acyclic acid being selected from the group consisting of
citric, lactic, tartaric, malic, glycolic, tannic and ascorbic
30 acids.

27. The method of claim 26, wherein said hydroxy acyclic
acid is citric acid.
35

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/01283

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :A01N 25/00 US CL :424/405 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 424/405 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 4,083,954 (TSUCHIYA ET AL.) 11 April 1978; see entire document.	1-27
Y	US, A, 3,867,546 (LECHEVIN ET AL.) 18 February 1975; see entire document.	1-27
Y	US, A, 4,581,378 (LAZAR ET AL.) 08 April 1986; see entire document.	1-27
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search 03 MARCH 1995	Date of mailing of the international search report 08 . 04 . 95	
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>Raj Bawa</i> RAJ BAWA, Ph.D. Telephone No. (703) 308-2351	